Complete the following problems. You must show your work to receive full credit. Show your answers to the correct number of significant figures with the correct units.

1. In the laboratory, you weigh out 0.114 grams of solid AlCl₃•6H₂O (molar mass 241.43 g/mol), dissolve it, and dilute it to a total volume of 100.0 mL to make solution A. You then transfer 3.00 mL of solution A into a 25.0 mL volumetric flask and dilute to the mark to make solution B. What is the molarity of aluminum ion in solution B? (9 pts.)

2. The 50.0 g Ca(OCl)₂ (molar mass 142.98 g/mol) and 275 mL of 3.00 M HCl are allowed to react in the reaction below. If 15.8 g of Cl₂ is produced, what is the percent yield for the reaction? (9 pts.)

 $Ca(OCI)_2(s) + 4HCI(aq) \rightarrow 2CI_2(g) + CaCI_2(aq) + 2H_2O(I)$

3. Billions of kilograms of urea (CO(NH₂)₂ molar mass = 60.06 g/mo)) are produced each year for use as fertilizer. The reaction involves combining ammonia (NH₃) with carbon dioxide to produce urea and water. What mass of carbon dioxide is required to convert 1.00 kg of ammonia to urea? What mass of urea is formed? (8 pts.)

Possibly Useful Information

| 1 | | | | | | | | | | | | | | | | | 18 |
|------------------------------|---------------------|----------------------|--------------------|---------------------|---------------------|---------------------|---------------------|------------------------|---------------------|---------------------|--------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
| 1A | | | | | | | | | | | | | | | | | 8A |
| 1 H | 2 | | | | | | | | | | | 13 | 14 | 15 | 16 | 17 | 2 He |
| 1.00794 | 2A | | | | | | | | | | | 3A | 4A | 5A | 6A | 7A | 4.00260 |
| 3 Li 6.941 | 4 Be 9.01218 | | | | | | | | | | | 5 B 10.811 | 6 C 12.011 | 7 N 14.0067 | 8 O 15.9994 | 9 F 18.9984 | 10 Ne 20.1797 |
| 11 Na 22.9898 | 12 Mg 24.3050 | 3 3B | 4 4B | 5 5B | 6 6B | 7 7B | 8 | 9 - _{8B} - | 10 | 11 1B | 12 2B | 13 Al 26.9815 | 14 Si 28.0855 | 15 P 30.9738 | 16 S 32.066 | 17 Cl 35.4527 | 18 Ar 39.948 |
| 19 K 39.0983 | 20 Ca 40.078 | 21 Sc 44.9559 | 22 Ti 47.88 | 23 V 50.9415 | 24 Cr 51.9961 | 25 Mn 54.9381 | 26 Fe 55.847 | 27 Co 58.9332 | 28 Ni 58.693 | 29 Cu 63.546 | 30 Zn 65.39 | 31 Ga 69.723 | 32 Ge 72.61 | 33 As 74.9216 | 34 Se 78.96 | 35 Br 79.904 | 36 Kr 83.80 |
| 37 Rb 85.4678 | 38 Sr 87.62 | 39 Y 88,9059 | 40 Zr 91,224 | 41 Nb 92,9064 | 42 Mo 95.94 | 43 Tc (98) | 44 Ru 101.07 | 45 Rh 102.906 | 46 Pd 106,42 | 47 Ag 107.868 | 48 Cd 112,411 | 49 In 114.818 | 50 Sn 118.710 | 51 Sb 121.757 | 52 Te 127.60 | 53 I 126.904 | 54 Xe 131.29 |
| 55 Cs 132.905 | 56 Ba 137,327 | 57 *La 138.906 | 72 Hf 178.49 | 73 Ta 180.948 | 74 W 183,84 | 75 Re 186.207 | 76 Os 190.23 | 77 Ir 192.22 | 78 Pt 195.08 | 79 Au 196,967 | 80 Hg 200,59 | 81 T1 204.383 | 82 Pb 207.2 | 83 Bi 208.980 | 84 Po (209) | 85 At (210) | 86 Rn (222) |
| 87 Fr (223) | 88 Ra 226.025 | 89 †Ac 227.028 | 104 Rf (261) | 105 Db (262) | 106 Sg (266) | 107 Bh (264) | 108 Hs (277) | 109 Mt (268) | 110 Ds (271) | 111 Rg (272) | | | | | | | |
| | | | | | | | | | | | | | | | | | |
| *Lanthanide series | | | | 58 Ce 140.115 | 59 Pr 140.908 | 60 Nd 144.24 | 61 Pm (145) | 62 Sm 150.36 | 63 Eu 151.965 | 64 Gd 157.25 | 65 Tb 158.925 | 66 Dy 162.50 | 67 Ho 164.930 | 68 Er 167.26 | 69 Tm 168.934 | 70 Yb 173.04 | 71 Lu 174.967 |
| [†] Actinide series | | | | 90 Th 232.038 | 91 Pa 231.036 | 92 U 238.029 | 93 Np 237.048 | 94 Pu (244) | 95 Am (243) | 96 Cm (247) | 97 Bk (247) | 98 Cf (251) | 99 Es (252) | 100 Fm (257) | 101 Md (258) | 102 No (259) | 103 Lr (262) |

Copyright © 2007 Pearson Prentice Hall, Inc.