

Chem 130
Exam 1, Ch 1-4
100 Points

Name _____
September 23, 2011

Please follow the instructions for each section of the exam. Show your work on all mathematical problems. Provide answers with the correct units and significant figures. Be concise in your answers to discussion questions.

Part 0: Warmup. 4 points each

- Thallium has two stable isotopes, ^{203}Tl and ^{205}Tl . Given that the atomic mass of thallium is 204.383 amu, which isotope must have the larger natural abundance?
 - ^{203}Tl
 - ^{205}Tl
 - Both have the same natural abundance.
 - Not enough information to make this determination.

Answer **B**
- Which of the following aspects of Dalton's atomic theory remains unchanged in our current understanding?
 - Atoms are indivisible.
 - All atoms of a particular element are identical.
 - Compounds are the result of a combination of two or more different kinds of atoms in fixed ratios.
 - None of the above.

Answer **C**

Part I: Complete all of problems 3-9

- Define the following using a maximum of two sentences for each definition. (8 points)
 - accuracy: **The proximity of a data point to the "true value"**
 - precision: **The reproducibility of a measurement or set of data.**

- Complete the following table. (12 points)

Symbol	$^{34}\text{S}^{2-}$	$^{40}\text{Ca}^{2+}$	^{58}Ni
# of protons	16	20	28
# of neutrons	16	20	30
# of electrons	18	18	28
Charge	-2	+2	0
Name	sulfide ion	calcium-40 ion	nickel-58

5. Name the following compounds or provide the correct formula for the given names. (18 pts)


- | | |
|--|---|
| a. diphosphorous tetrafluoride | <u>P₂F₄</u> |
| b. Al ₂ (CO ₃) ₃ | <u>aluminum carbonate</u> |
| c. Cr(PO ₄) ₂ | <u>chromium (VI) phosphate</u> |
| d. iron (III) sulfate | <u>Fe₂(SO₄)₃</u> |
| e. calcium chloride | <u>CaCl₂</u> |
| f. N ₂ O ₅ | <u>dinitrogen pentoxide</u> |

6. How many ²⁰⁴Pb atoms are in a piece of lead weighing 215 mg? The percent natural abundance of lead is 1.4%. (8 points)


$$0.215 \text{ g-Pb} \times \frac{1 \text{ mol-Pb}}{207.2 \text{ g-Pb}} \times \frac{6.02 \times 10^{23} \text{ atoms-Pb}}{1 \text{ mol-Pb}} \times \frac{1.4 \text{ atoms } ^{204}\text{Pb}}{100 \text{ atoms-Pb}} = 8.8 \times 10^{18} \text{ } ^{204}\text{Pb atoms}$$

7. Write balanced reactions, specifying the state for all reactants and products. (8 points)

- a. Aqueous copper (I) sulfate reacts with aqueous barium iodide to produce solid barium sulfate and aqueous copper (I) iodide.



- b. Aqueous sodium carbonate reacts with gaseous nitrogen monoxide and oxygen gas to produce aqueous sodium nitrate and carbon monoxide gas.



8. A solution consisting of 8.50% acetone and 91.50% water by mass has a density of 0.9867 g/mL. What mass of acetone, in kg, is present in 7.50 L of the solution? (8 pts)

$$7500 \text{ mL} \times \frac{0.9867 \text{ g-soln}}{\text{mL}} \times \frac{8.50 \text{ g acetone}}{100 \text{ g-soln}} \times \frac{1 \text{ kg}}{10^3 \text{ g}} = 0.629 \text{ kg acetone}$$

Part II. Answer three (3) of problems 9-12. Clearly mark the problem you do not want graded. 10 points each.

9. Silicon has three stable isotopes, ^{28}Si , ^{29}Si , and ^{30}Si with masses of 27.98 amu, 28.98 amu, and 29.77 amu, respectively. If the natural abundance of ^{28}Si is 92.23%, what are the percent abundances of the other two isotopes?

The total abundance of ^{29}Si and ^{30}Si must be: $100 - 92.23 = 7.77\%$

So: $f_{29} + f_{30} = 0.0777$ where f_x is the fractional abundance of the isotope with mass # x.


And: $(0.9223 \times 27.98) + 28.98f_{29} + 29.77f_{30} = 28.0855$ (This is our definition of atomic mass)

Now we need to find f_{29} and f_{30} :

$$\begin{aligned} f_{30} &= 0.0777 - f_{29} \\ 28.98f_{29} + 29.77f_{30} &= 28.0855 - (0.9223 \times 27.98) = 2.2795 \\ 28.98f_{29} + 29.77(0.0777 - f_{29}) &= 2.2795 \\ 28.98f_{29} - 29.77f_{29} &= 2.2795 - (29.77 \times 0.0777) = -0.03363 \\ -0.79f_{29} &= -0.03363 \\ f_{29} &= 0.04257 \\ \text{So, } f_{30} &= 0.0777 - 0.04257 = 0.03513 \end{aligned}$$

So, the percent abundance for ^{29}Si is 4.26% and the percent abundance for ^{30}Si is 3.51%

10. Ammonia can be generated by heating together the solids $\text{Ca}(\text{OH})_2$ and NH_4Cl . CaCl_2 and water are also formed. How many grams of NH_3 will form if 33.0 grams each of NH_4Cl and $\text{Ca}(\text{OH})_2$ are heated? (molar masses (g/mol): $\text{NH}_4\text{Cl} = 53.4912$, $\text{NH}_3 = 17.03056$, $\text{Ca}(\text{OH})_2 = 74.093$, $\text{CaCl}_2 = 110.983$, water = 18.0153)



If NH_4Cl is the limiting reactant, how many grams of ammonium could be produced?

$$33.0 \text{ g } \text{NH}_4\text{Cl} \times \frac{1 \text{ mol } \text{NH}_4\text{Cl}}{53.4912 \text{ g } \text{NH}_4\text{Cl}} \times \frac{2 \text{ mol } \text{NH}_3}{2 \text{ mol } \text{NH}_4\text{Cl}} \times \frac{17.03056 \text{ g } \text{NH}_3}{1 \text{ mol } \text{NH}_3} = 10.51 \text{ g } \text{NH}_3$$

If $\text{Ca}(\text{OH})_2$ is the limiting reactant, how many grams of ammonium could be produced?

$$33.0 \text{ g } \text{Ca}(\text{OH})_2 \times \frac{1 \text{ mol } \text{Ca}(\text{OH})_2}{74.093 \text{ g } \text{Ca}(\text{OH})_2} \times \frac{2 \text{ mol } \text{NH}_3}{1 \text{ mol } \text{Ca}(\text{OH})_2} \times \frac{17.03056 \text{ g } \text{NH}_3}{1 \text{ mol } \text{NH}_3} = 15.17 \text{ g } \text{NH}_3$$

Therefore, ammonium chloride must be the limiting reagent, and a maximum of 10.5 grams of ammonia could be produced.

11. Iron ore is impure Fe_2O_3 . When Fe_2O_3 is heated with carbon, metallic iron and carbon monoxide gas are formed. From a sample of ore weighing 938 g, 532 g of pure iron is obtained. What is the percent Fe_2O_3 , by mass, in the original ore sample? (molar masses (g/mol): $\text{Fe}_2\text{O}_3 = 159.6922$, carbon monoxide = 28.010)



$$532 \text{ g Fe} \times \frac{1 \text{ mol Fe}}{55.847 \text{ g Fe}} \times \frac{1 \text{ mol Fe}_2\text{O}_3}{2 \text{ mol Fe}} \times \frac{159.6922 \text{ g Fe}_2\text{O}_3}{1 \text{ mol Fe}_2\text{O}_3} = 760.6 \text{ g Fe}_2\text{O}_3$$

Therefore, 761 grams of Fe_2O_3 must have been present in the original sample. In terms of percent, this corresponds to:

$$\frac{761 \text{ g Fe}_2\text{O}_3}{938 \text{ g ore}} \times 100\% = 81.1\% \text{ Fe}_2\text{O}_3$$

12. One of the reasons that methamphetamine is such a problem is that it is a relatively small molecule that is fairly easy to synthesize. A molecule of methamphetamine contains only carbon, hydrogen, and nitrogen and has a molar mass of 149.2 g/mol. If methamphetamine is 80.48% C and 9.39% N by mass, what is its molecular formula?

So, the percent H must be: $100 - 80.48 - 9.39 = 10.13\% \text{H}$

There are several ways to solve this problem. Here is one:

$$\frac{149.2 \text{ g-meth}}{1 \text{ mol meth}} \times \frac{80.48 \text{ g-C}}{100 \text{ g-meth}} \times \frac{1 \text{ mol C}}{12.011 \text{ g-C}} = \frac{9.997 \text{ mol C}}{1 \text{ mol meth}}$$

$$\frac{149.2 \text{ g-meth}}{1 \text{ mol meth}} \times \frac{9.39 \text{ g-N}}{100 \text{ g-meth}} \times \frac{1 \text{ mol N}}{14.0067 \text{ g-N}} = \frac{1.000 \text{ mol N}}{1 \text{ mol meth}}$$

$$\frac{149.2 \text{ g-meth}}{1 \text{ mol meth}} \times \frac{10.13 \text{ g-H}}{100 \text{ g-meth}} \times \frac{1 \text{ mol H}}{1.00794 \text{ g-H}} = \frac{14.995 \text{ mol H}}{1 \text{ mol meth}}$$

So, the likely formula is $\text{C}_{9.997}\text{H}_{14.995}\text{N}_{1.000}$ or, $\text{C}_{10}\text{H}_{15}\text{N}$. Is this really the molecular formula?

Check the molar mass:

$$10(12.011) + 15(1.00794) + 14.0067 = 149.23$$

Therefore, $\text{C}_{10}\text{H}_{15}\text{N}$ must be the molecular formula!

Form A
Possibly Useful Information

D = m/v	$N_a = 6.02214 \times 10^{23} \text{ mol}^{-1}$
---------	---

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1A	2A											3A	4A	5A	6A	7A	8A
1 H 1.00794																	2 He 4.00260
	3 Li 6.941	4 Be 9.01218										5 B 10.811	6 C 12.011	7 N 14.0067	8 O 15.9994	9 F 18.9984	10 Ne 20.1797
11 Na 22.9898	12 Mg 24.3050		3 B	4 C	5 N	6 O	7 F	8 Ne	9 Na	10 Mg	11 Al 26.9815	12 Si 28.0855	13 P 30.9738	14 S 32.066	15 Cl 35.4527	16 Ar 39.948	
19 K 39.0983	20 Ca 40.078	21 Sc 44.9559	22 Ti 47.88	23 V 50.9415	24 Cr 51.9961	25 Mn 54.9381	26 Fe 55.847	27 Co 58.9332	28 Ni 58.693	29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.9216	34 Se 78.96	35 Br 79.904	36 Kr 83.80
37 Rb 85.4678	38 Sr 87.62	39 Y 88.9059	40 Zr 91.224	41 Nb 92.9064	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.906	46 Pd 106.42	47 Ag 107.868	48 Cd 112.411	49 In 114.818	50 Sn 118.710	51 Sb 121.757	52 Te 127.60	53 I 126.904	54 Xe 131.29
55 Cs 132.905	56 Ba 137.327	57 *La 138.906	72 Hf 178.49	73 Ta 180.948	74 W 183.84	75 Re 186.207	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.967	80 Hg 200.59	81 Tl 204.383	82 Pb 207.2	83 Bi 208.980	84 Po (209)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra 226.025	89 *Ac 227.028	104 Rf (261)	105 Db (262)	106 Sg (266)	107 Bh (264)	108 Hs (277)	109 Mt (268)	110 Ds (271)	111 Rg (272)							

*Lanthanide series	58 Ce 140.115	59 Pr 140.908	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.965	64 Gd 157.25	65 Tb 158.925	66 Dy 162.50	67 Ho 164.930	68 Er 167.26	69 Tm 168.934	70 Yb 173.04	71 Lu 174.967
+ Actinide series	90 Th 232.038	91 Pa 231.036	92 U 238.029	93 Np 237.048	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (262)

Copyright © 2007 Pearson Prentice Hall, Inc.