

FORM B

Chem 130
Exam 1, Ch 5-6
100 Points

Name _____
October 21, 2011

Please follow the instructions for each section of the exam. Show your work on all mathematical problems. Provide answers with the correct units and significant figures. Be concise in your answers to discussion questions. Point values are in parentheses by each problem.

Part 0: Warmup. 4 points each

1. Choose the INCORRECT statement:
- a. Most molecular compounds are either non-electrolytes or weak electrolytes.
 - b. An acid produces hydride ions in solution. Answer _____
 - c. Net ionic equations include only the actual participants in the reaction.
 - d. Most ionic compounds are strong electrolytes.
2. If someone were to light a cigar at one end of a closed room, persons on the other end of the room might soon perceive an odor due to gaseous emissions from the cigar. Such a phenomenon is an example of:
- a. ideality
 - b. effusion
 - c. dissolution
 - d. diffusion Answer _____

Part I: Complete all of problems 3-8

3. Compare the following using a maximum of three sentences for each pair of terms.
- a. **strong electrolyte** versus **weak electrolyte**. (5)

 - b. **ideal gas** versus **non-ideal (or real) gas**: (5)
4. It is possible to use precipitation reactions to separate ions in solution by removing target ions as insoluble salts. Propose an approach to separate Fe^{2+} from Ca^{2+} using precipitation reactions. Include balanced reactions (indicating states of products and reactants). (10)

FORM B

5. Answer the following based on the reaction: $\text{SO}_3^{2-} + \text{MnO}_4^- \rightarrow \text{SO}_4^{2-} + \text{Mn}^{2+}$
- What is the oxidation state of manganese in the permanganate ion? _____ (2)
 - What is the oxidation state of sulfur in the sulfate ion? _____ (2)
 - Balance the reaction in acidic aqueous solution. (8)
6. A 1.27 g sample of an oxide of nitrogen, believed to be either NO or N_2O , occupies a volume of 0.728 L at 25°C and 737 mm Hg. Which oxide is it? (10)

FORM B

7. A 0.755 gram sample of solid magnesium hydroxide is added to 125 mL of a 0.555 M solution of nitric acid. Will the resulting solution be acidic, basic, or neutral? Justify your answer. (10)

8. How does the kinetic-molecular theory of gases help explain why a helium-filled balloon shrinks if it is taken outside on a cold winter day? (10)

FORM B

Part II. Answer two (2) of problems 9-12. Clearly mark the problems you do not want graded. 15 points each.

9. Write balanced overall reactions and net ionic equations for each of the following: Indicate the state (*s*, *l*, *g*, *aq*) of each of the reactants and products.

a. Aqueous lead (II) nitrate is mixed with aqueous lithium hydroxide

Balanced Reaction: (5)

Net Ionic Equation: (2)

b. Aqueous sulfuric acid is mixed with aqueous sodium bicarbonate

Balanced Reaction: (5)

Net Ionic Equation: (2)


10. Sodium dithionite, $\text{Na}_2\text{S}_2\text{O}_4$, is an important reducing agent. One interesting use is in the purification of wastewater by the reduction of chromate ion with $\text{S}_2\text{O}_4^{2-}$ in basic solution to form insoluble chromium (III) hydroxide, with sulfite ion produced as another product.

a. Write the balanced reaction for the process occurring in basic solution. (10)


b. What mass of $\text{Na}_2\text{S}_2\text{O}_4$ is consumed in a reaction with 100.0 L of wastewater having $[\text{CrO}_4^{2-}] = 0.0166 \text{ M}$? (5)

FORM B

11. Birmabright is a metal alloy consisting of aluminum, magnesium, and manganese. A 0.273 g sample of Birmabright is dissolved in an excess of hydrochloric acid, producing hydrogen gas as shown in the balanced reaction below. If 355 mL of hydrogen is collected over water at a temperature of 25°C and pressure of 755 mm Hg, what is the mass percent of aluminum in Birmabright?



12. Nitroglycerine ($\text{C}_3\text{H}_5\text{N}_3\text{O}_9$, molar mass = 227.088 g/mol) is a contact explosive that rapidly decomposes via the reaction below and releases a large quantity of heat and gas. Assume 10.0 grams of nitroglycerine decomposes in a 2.0 L soda bottle and instantaneously generates a temperature of 5230K.



- a. What will the pressure be inside the bottle once the reaction is complete? (5)
- b. What is the partial pressure of water when the reaction is complete? (5)
- c. What volume would the gas mixture produced occupy at STP? (5)

FORM B
Possibly Useful Information

$R = 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1}$	STP = 1 atm, 0°C $K = 273.15 + ^\circ\text{C}$
1 atmosphere = 760 Torr = 760 mm Hg	$\left(P + \frac{n^2a}{V^2}\right)(V - bn) = nRT$
$P_{\text{total}} = n_{\text{total}}RT/V$	$P_A = X_A P_{\text{total}}$
$N_a = 6.02214 \times 10^{23} \text{ mol}^{-1}$	$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$

Molar Masses	
Compound	Molar Mass (g/mol)
aluminum chloride	133.340
carbon dioxide	44.010
chromium (III) hydroxide	103.018
hydrochloric acid	36.461
hydrogen gas	2.016
magnesium hydroxide	58.320
nitric acid	63.013
nitrogen gas	28.0135
nitrogen monoxide	30.006
nitroglycerine	227.088
oxygen gas	31.999
sodium carbonate	105.989
sodium dithionite	174.109
sodium nitrite	68.995
water	18.015

Vapor Pressure of Water at Various Temperatures

Temperature (°C)	Vapor Pressure (mmHg)
15.0	12.79
17.0	14.53
19.0	16.48
21.0	18.65
23.0	21.07
25.0	23.76
30.0	31.82
50.0	92.51

Copyright © 2007 Pearson Prentice Hall, Inc.

TABLE 5.3 Some Common Gas-Forming Reactions

Ion	Reaction
HSO_3^-	$\text{HSO}_3^- + \text{H}^+ \longrightarrow \text{SO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
SO_3^{2-}	$\text{SO}_3^{2-} + 2 \text{H}^+ \longrightarrow \text{SO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
HCO_3^-	$\text{HCO}_3^- + \text{H}^+ \longrightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
CO_3^{2-}	$\text{CO}_3^{2-} + 2 \text{H}^+ \longrightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
S^{2-}	$\text{S}^{2-} + 2 \text{H}^+ \longrightarrow \text{H}_2\text{S}(\text{g})$
NH_4^+	$\text{NH}_4^+ + \text{OH}^- \longrightarrow \text{NH}_3(\text{g}) + \text{H}_2\text{O}(\text{l})$

Copyright © 2007 Pearson Prentice Hall, Inc.

TABLE 5.1 Solubility Guidelines for Common Ionic Solids

Follow the lower-numbered guideline when two guidelines are in conflict. This leads to the correct prediction in most cases.

- Salts of group 1 cations (with some exceptions for Li^+) and the NH_4^+ cation are soluble.
- Nitrates, acetates, and perchlorates are soluble.
- Salts of silver, lead, and mercury(I) are insoluble.
- Chlorides, bromides, and iodides are soluble.
- Carbonates, phosphates, sulfides, oxides, and hydroxides are insoluble (sulfides of group 2 cations and hydroxides of Ca^{2+} , Sr^{2+} , and Ba^{2+} are slightly soluble).
- Sulfates are soluble except for those of calcium, strontium, and barium.

FORM B

1	2											18	19															
1A	2A											8A	8B															
¹ H 1.00794																				² He 4.00260								
3 Li 6.941	4 Be 9.01218																				5 B 10.811	6 C 12.011	7 N 14.0067	8 O 15.9994	9 F 18.9984	10 Ne 20.1797		
11 Na 22.9898	12 Mg 24.3050	3 3B	4 4B	5 5B	6 6B	7 7B	8	9	10	11	12	13 Al 26.9815	14 Si 28.0855	15 P 30.9738	16 S 32.066	17 Cl 35.4527	18 Ar 39.948											
19 K 39.0983	20 Ca 40.078	21 Sc 44.9559	22 Ti 47.88	23 V 50.9415	24 Cr 51.9961	25 Mn 54.9381	26 Fe 55.847	27 Co 58.9332	28 Ni 58.693	29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.9216	34 Se 78.96	35 Br 79.904	36 Kr 83.80											
37 Rb 85.4678	38 Sr 87.62	39 Y 88.9059	40 Zr 91.224	41 Nb 92.9064	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.906	46 Pd 106.42	47 Ag 107.868	48 Cd 112.411	49 In 114.818	50 Sn 118.710	51 Sb 121.757	52 Te 127.60	53 I 126.904	54 Xe 131.29											
55 Cs 132.905	56 Ba 137.327	57 *La 138.906	72 Hf 178.49	73 Ta 180.948	74 W 183.84	75 Re 186.207	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.967	80 Hg 200.59	81 Tl 204.383	82 Pb 207.2	83 Bi 208.980	84 Po (209)	85 At (210)	86 Rn (222)											
87 Fr (223)	88 Ra 226.025	89 †Ac 227.028	104 Rf (261)	105 Db (262)	106 Sg (266)	107 Bh (264)	108 Hs (277)	109 Mt (268)	110 Ds (271)	111 Rg (272)																		
*Lanthanide series		58 Ce 140.115	59 Pr 140.908	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.965	64 Gd 157.25	65 Tb 158.925	66 Dy 162.50	67 Ho 164.930	68 Er 167.26	69 Tm 168.934	70 Yb 173.04	71 Lu 174.967													
†Actinide series		90 Th 232.038	91 Pa 231.036	92 U 238.029	93 Np 237.048	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (262)													