Chemistry 222	
Spring 2014	
Exam 2: Chapters	ô-7

Name	
	80 Points

Complete problem 1 and four of problems 2-6. CLEARLY mark the problem you do not want graded. You must show your work to receive credit for problems requiring math. Report your answers with the appropriate number of significant figures.

Bonus (4 points):

One day last week, Dr. Lamp told those in class a pH at 8:30 AM as he started class and said that the pH would be an answer to one of the exam 2 questions. What was that pH?_____

You MUST complete problem 1. (16 pts.)

1. When ammonium sulfate dissolves, both the anion and the cation can participate in acid-base equilibria. Considering all the equilibria active in solution, write enough equations so that you could solve for the concentration of each species in a solution that is saturated with ammonium sulfate that also contains 0.10 M sodium nitrate. You must write the charge balance expression and at least one mass balance. Identify all unknowns and write enough explicit, independent equations so that only algebra remains to solve for the unknowns. A numerical answer is not necessary. Do not consider activities.

 $\begin{array}{ll} (NH_4)_2SO_4(s) & K_{sp} = 276 \\ NH_4^+ & K_a = 5.7 \times 10^{-10} \\ SO_4^{2-} & K_b = 9.8 \times 10^{-13} \end{array}$

Do four of problems 2-6. Clearly mark the problem you do not want graded. (16 pts. ea.)

2. A saturated solution of BaSO₄ ($K_{sp} = 1.1 \times 10^{-10}$) that originally held a volume of 1.00 L is allowed to evaporate until the solution volume is 0.500 L. How does the new concentration of Ba²⁺ compare to the concentration in the original solution? Clearly justify your response, using calculations were appropriate. *Do not consider activities*.

3. Is it possible to perform a 99.99 % complete separation of barium and silver by precipitation with sulfate if both Ba²⁺ and Ag⁺ are present initially at 0.020 M? Justify your decision. *Do not consider activities*.

$$\begin{array}{ll} BaSO_4 & K_{sp} = 1.1 \ x \ 10^{\text{-}10} \\ Ag_2SO_4 & K_{sp} = 1.4 \ x \ 10^{\text{-}5} \end{array}$$

4. Consider the table of activity coefficients on the last page of this exam. As you move from left to right across any row on the table, the values for activity coefficient decrease. As you move down in a given column, the activity coefficient also decreases. Clearly describe the phenomena that cause these trends. Do not simply point out the trends; you must explain why the trends exist. No calculations are necessary.



6.	Ethylamine (CH $_3$ CH $_2$ NH $_2$) is a monobasic weak base with a pK $_b$ of 3.33. Calculate the pH of a solution prepared by mixing 20.0 mL of 0.010 M HCl with 50.0 mL of 0.046 M ethylamine (CH $_3$ CH $_2$ NH $_2$) and diluting the resulting solution to 100.0 mL. <i>Do not consider activities</i> .						

Possibly Useful Information

$K_a K_b = K_W = 1.0 \times 10^{-14}$	Don't eat the yellow snow.
$-\log\gamma = \frac{0.51z^2\sqrt{\mu}}{1+\alpha\sqrt{\mu}/305}$	$\mu = \frac{1}{2} \sum_{i} c_{i} z_{i}^{2}$
$\Delta G = \Delta H - T\Delta S = -RTInK$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
$H_2O \Rightarrow H^+ + OH^- K_w = 1.00x10^{-14}$	pH = -log[H ⁺]

Activity coefficients for aqueous solutions at 25°C

	Ion size	Ionic strength (μ, M)				
Ion	(α, pm)	0.001	0.005	0.01	0.05	0.1
$CHARGE = \pm 1$						
H ⁺	900	0.967	0.933	0.914	0.86	0.83
$(C_6H_5)_2CHCO_2^-, (C_3H_7)_4N^+$	800	0.966	0.931	0.912	0.85	0.82
$(O_2N)_3C_6H_2O^-, (C_3H_7)_3NH^+, CH_3OC_6H_4CO_2^-$	700	0.965	0.930	0.909	0.845	0.81
Li ⁺ , C ₆ H ₅ CO ₂ , HOC ₆ H ₄ CO ₂ , ClC ₆ H ₄ CO ₂ , C ₆ H ₅ CH ₂ CO ₂ ,						
$CH_2 = CHCH_2CO_2^-, (CH_3)_2CHCH_2CO_2^-, (CH_3CH_2)_4N^+, (C_3H_7)_2NH_2^+$	600	0.965	0.929	0.907	0.835	0.80
Cl ₂ CHCO ₂ -, Cl ₃ CCO ₂ -, (CH ₃ CH ₂) ₃ NH ⁺ , (C ₃ H ₇)NH ₃ ⁺	500	0.964	0.928	0.904	0.83	0.79
Na^{+} , $CdCl^{+}$, ClO_{2}^{-} , IO_{3}^{-} , HCO_{3}^{-} , $H_{2}PO_{4}^{-}$, HSO_{3}^{-} , $H_{2}AsO_{4}^{-}$,						
$Co(NH_3)_4(NO_2)_2^+$, $CH_3CO_2^-$, $CICH_2CO_2^-$, $(CH_3)_4N^+$,						
$(CH_3CH_2)_2NH_2^+, H_2NCH_2CO_2^-$	450	0.964	0.928	0.902	0.82	0.775
+H ₃ NCH ₂ CO ₂ H, (CH ₃) ₃ NH ⁺ , CH ₃ CH ₂ NH ₃ ⁺	400	0.964	0.927	0.901	0.815	0.77
OH ⁻ , F ⁻ , SCN ⁻ , OCN ⁻ , HS ⁻ , ClO ₃ ⁻ , ClO ₄ ⁻ , BrO ₃ ⁻ , IO ₄ ⁻ , MnO ₄ ⁻ ,						
HCO ₂ ⁻ , H ₂ citrate ⁻ , CH ₃ NH ₃ ⁺ , (CH ₃) ₂ NH ₂ ⁺	350	0.964	0.926	0.900	0.81	0.76
K ⁺ , Cl ⁻ , Br ⁻ , I ⁻ , CN ⁻ , NO ₂ , NO ₃	300	0.964	0.925	0.899	0.805	0.755
$Rb^+, Cs^+, NH_4^+, TI^+, Ag^+$	250	0.964	0.924	0.898	0.80	0.75
$CHARGE = \pm 2$						
Mg^{2+}, Be^{2+}	800	0.872	0.755	0.69	0.52	0.45
$CH_2(CH_2CH_2CO_2^-)_2$, $(CH_2CH_2CH_2CO_2^-)_2$	700	0.872	0.755	0.685	0.50	0.425
Ca^{2+} , Cu^{2+} , Zn^{2+} , Sn^{2+} , Mn^{2+} , Fe^{2+} , Ni^{2+} , Co^{2+} , $C_6H_4(CO_2^-)_2$,						
H ₂ C(CH ₂ CO ₂ ⁻) ₂ , (CH ₂ CH ₂ CO ₂ ⁻) ₂	600	0.870	0.749	0.675	0.485	0.405
Sr^{2+} , Ba^{2+} , Cd^{2+} , Hg^{2+} , S^{2-} , $S_2O_4^{2-}$, WO_4^{2-} , $H_2C(CO_2^-)_2$, $(CH_2CO_2^-)_2$,						
$(CHOHCO_2^-)_2$	500	0.868	0.744	0.67	0.465	0.38
Pb^{2+} , CO_3^{2-} , SO_3^{2-} , MoO_4^{2-} , $Co(NH_3)_5Cl^{2+}$, $Fe(CN)_5NO^{2-}$, $C_2O_4^{2-}$,						
Hcitrate ²⁻	450	0.867	0.742	0.665	0.455	0.37
$\mathrm{Hg_{2}^{2+}}, \mathrm{SO_{4}^{2-}}, \mathrm{S_{2}O_{3}^{2-}}, \mathrm{S_{2}O_{6}^{2-}}, \mathrm{S_{2}O_{8}^{2-}}, \mathrm{SeO_{4}^{2-}}, \mathrm{CrO_{4}^{2-}}, \mathrm{HPO_{4}^{2-}}$	400	0.867	0.740	0.660	0.445	0.355
$CHARGE = \pm 3$						
Al^{3+} , Fe^{3+} , Cr^{3+} , Sc^{3+} , Y^{3+} , In^{3+} , lanthanides ^a	900	0.738	0.54	0.445	0.245	0.18
citrate ³⁻	500	0.728	0.51	0.405	0.18	0.115
PO_4^{3-} , $Fe(CN)_6^{3-}$, $Cr(NH_3)_6^{3+}$, $Co(NH_3)_6^{3+}$, $Co(NH_3)_5H_2O^{3+}$	400	0.725	0.505	0.395	0.16	0.095
$Charge = \pm 4$						
$Th^{4+}, Zr^{4+}, Ce^{4+}, Sn^{4+}$	1 100	0.588	0.35	0.255	0.10	0.065
Fe(CN) ₆ ⁴⁻	500	0.57	0.31	0.20	0.048	0.021
	0 10.02					

a. Lanthanides are elements 57–71 in the periodic table. SOURCE: J. Kielland, J. Am. Chem. Soc. 1937, 59, 1675.

PERIODIC CHART OF THE ELEMENTS IB IIB IIIA IA IIA IIIB IYB YB YIB YIIB YIII IVA VA VIA VIIA GASES 1 He H 1.00797 1.00797 Li 10 4 9 В C Be Ν O Ne 10.811 12.0112 18.998 9.0122 20.183 11 12 17 18 13 Si S 32.064 Р CI ΑI Na ∣Mg Ar 30.9738 28.086 35.453 39.948 22.9898 24.3T 26.981 20 26 35 19 31 32 33 34 36 K Ca Sc 44.956 Zn 65.37 Se 78.96 Co Kr 83.80 ٧ Cr Mn Fe Ni Τi **Ga** As Br Ge 58.9332 50.942 51.996 54.9380 55.847 74.9216 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 **Sb** Sr 87.62 Cd Sn 126.904 Xe Ŗb **Ru** Rh Ag 107.870 **Zr** Тс PdTe Nb Μo In 106.4 88.905 85.47 92.906 95.94 (99) 102.905 114.82 55 56 ***57** 74 81 72 73 75 76 77 78 79 80 82 83 84 85 86 Cs Ba Hg 200.59 **At** (210) **La** Po Hf Re Os Pt ΤI Pb Bi Rn Ta Ir 180.948 183.85 190.2 192.2 195.09 196.967 204.37 207.19 208.980 (210) (222) 87 88 **±89** 104 105 106 107 108 109 110 112 Rf Db Sg Bh Hs ? ? Mt ? Fr |Ra|Ac (272) (277) (271) (227) (226) Numbers in parenthesis are mass numbers of most stable or most *Lanthanide Series 58 59 60 61 62 63 64 common isotope. Ce Pr Nd Pm Sm Eu Gd Tb Dy Ho Er Tm Tm|Yb Lu Atomic weights corrected to conform to the 1963 values of the Commission on Atomic Weights. **†** Actinide Series 90 91 95 96 99 100 101 The group designations used here are the former Chemical Abstract Service numbers. Np Pu Am Cm Bk (247) (247) Es Fm Md No (256) (256) Th Pa Cf U

(249)

(257)

232.038 (231) 238.03