CHEM 131 Quiz 8 – Due in class November 20, 2019

Complete the following problems. Write your final answers in the blanks provided. You must show your work to receive full credit. Show your answers to the correct number of significant figures with the correct units.

Name _____

Rules for this take-home quiz.

DO NOT OPEN THE QUIZ UNTIL YOU ARE READY TO TAKE IT!

- You may allocate a maximum of **50 continuous minutes** for this quiz, split in to two 25-minute segments.
- For the first 25-minute segment, you will take the quiz using only the materials on these pages, a calculator and a **<u>pencil</u>**. Treat this time as though you were taking the quiz in the classroom. You may not use your book, notes, electronic sources or anyone else to help. Record the start and end of the first 25 minutes below.
- For the second 25 minutes, you may use your book, notes or electronic resources to make any corrections to your work. **Make these corrections in blue or red pen.** You **MAY NOT** ask anyone else for help. Record the end of the second 25 minute block below.
- Once you have completed the quiz, sign below to affirm that the quiz was taken following the rules above. This signature is your pledge that the quiz was completed in an ethical manner!

Start time:	End of 1 st 25 minutes:	End of 2 nd 25 minutes:
Signature		Date

Possibly Useful Information

$E^{o}_{cell} = E^{o}_{cathode} - E^{o}_{anode}$	2 303PT		$R = 8.314 L atm mol^{-1} K^{-1}$
or	$E = E^{\circ} - \frac{2.505 \text{ KT}}{1000} \log Q$	$\Delta G = -nFE$	$F = 96485 \text{ C mol}^{-1}$
$E^{o}_{cell} = E^{o}_{cathode} - E^{o}_{anode}$	nF		$1 \text{ A} = 1 \text{ C sec}^{-1}$

IA																	VIIIA
1A	1					Dori	odic 1	Tabla /	of the		onte						8A
Ľω.						ren	ouic	able		Eleli	ients						ί μ _α
	2											13	14	15	16	17	пе
1.008	24											34	44	54	64	74	4.003
3	4											5	6	7	8	9	10
11	Re											R	C	N	0	F	No
Lithium	Beryllium											Boron	Carbon	Nitrogen	Oxygen	Fluorine	Neon
6.941	9.012											10.811	12.011	14.007	15.999	18.998	20.180
11	12											13	14	15	16	17	18
Na	Ma	3	4	5	6	7	8	9	10	11	12		Si	P	S	C	Ar
Sodium	Magnesium	шв	IVB	VB	VIB	VIIB		— vIII —		IB	ШΒ	Aluminum	Silicon	Phosphorus	Sulfur	Chlorine	Argon
22.990	24.305	3B	4B	5B	6B	7B	*	8	*	1B	2B	26.982	28.086	30.974	32.066	35.453	39.948
19	20	²¹ C -	²² - :	23	²⁴ C	25 B.4	26 F -	" c -	28	29	30	³¹	³²	33	³⁴ C -	35 D	30
K	Ca	SC		V	Cr	ivin	ге	CO	NI	Cu	Zn	Ga	Ge	AS	Se	Br	Kr
Potassium 39.098	Calcium 40.078	Scandium 44.956	Titanium 47.867	Vanadium 50.942	Chromium 51.996	Manganese 54.938	Iron 55.845	Cobalt 58.933	Nickel 58.693	Copper 63.546	Zinc 65.38	Gallium 69.723	Germanium 72.631	Arsenic 74.922	Selenium 78.971	Bromine 79.904	Krypton 83.798
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rh	Sr	V	7r	Nh	Mo	Тс	Ru	Rh	Pd	Δα	Cd	In	Sn	Sh	Te	T	Xe
Rubidium	Strontium	Yttrium	Zirconium	Niobium	Molybdenum	Technetium	Ruthenium	Rhodium	Palladium	Silver	Cadmium	Indium	Tin	Antimony	Tellurium	Iodine	Xenon
85.468	87.62	88.906	91.224	92.906	95.95	98.907	101.07	102.906	106.42	107.868	112.414	114.818	118.711	121.760	127.6	126.904	131.294
55	56_	57-71	72	73	74	75	76	77 _	78	79	80	81	82	83	84_	85	86
Cs	Ba		Hf	Та	W	Re	Os	Ir	Pt	Au	Hg	TI	Pb	Bi	Po	At	Rn
Cesium	Barium		Hafnium	Tantalum	Tungsten	Rhenium	Osmium	Iridium	Platinum	Gold	Mercury	Thallium	Lead	Bismuth	Polonium	Astatine	Radon
132.905	00	90-102	104	100.946	105.04	100.207	190.23	100	110	196.967	112	204.363	114	115	116	117	110
°' E =	° Da	89-105	Df	Dh.	С.я.	Ph		N/+	Da	Da	"Cn	NIL	EI	Ma	1.	"To	
Fr	Rđ		KI	DD	5 g	DU	ПS	IVIT	DS	ĸg	Cn	INN	ГІ	IVIC	LV	IS	Ug
223.020	226.025		[261]	[262]	[266]	[264]	[269]	[278]	[281]	[280]	[285]	[286]	[289]	[289]	[293]	[294]	[294]
		57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	
	Lanth	anide	.a C	Ce F	Pr N	d P	m S	m E	u l C	id 1	「b D)v ⊦	lo E	Er T	ím 🛛 ۱	/b L	.u 🗌
	Ser	Lant	hanum Ce	rium Prasec	dymium Neod	mium Prom	ethium Sam	narium Euro	opium Gad	olinium Te	rbium Dysp	rosium Ho	lmium Er	bium Th	ulium Ytte	erbium Lut	etium
		13	8.905 14	0.116 14	1.908 144	243 14	1913 19	0.36 151	1.964 15	07.25 15	8.925 162		4.930 16	7.259 16	8.934 17	3.055 17	4.967
	Activ	nide 89	90		92	93	94 F	95 A		⁹⁷ F	98	יב ⁹⁹ ד		101		103	
	Ser	ies F			al	ע ו	ір і Р	'u A	mC	m	SK (J I	:S F	m N		10	Lr
		Act 22	inium Th 7.028 23	2.038 Prota 2.038 23	ctinium Uran 1.036 238	029 Nept 029 23	unium Plut 7.048 24	onium Ame 4.064 243	ricium Cu 3.061 24	rium Ber 7.070 24	kelium Califo 7.070 251	Einst 1.080	teinium Fer 254] 25	mium Mend 7.095 2	Selevium No 58.1 25	belium Lawr 9.101 [encium 262]

© 2017 Todd Helmenstine sciencenotes.org

TABLE 20.1	Some	Selected Sta	ndard	Electrode	(Reduction)
Potentials at	25 °C				

Reduction Half-Reaction	<i>E</i> °, V					
Acidic solution						
$F_2(g) + 2 e^- \longrightarrow 2 F^-(aq)$	+2.866					
$O_3(g) + 2 H^+(aq) + 2 e^- \longrightarrow O_2(g) + H_2O(l)$	+2.075					
$S_2O_8^{2-}(aq) + 2e^- \longrightarrow 2SO_4^{2-}(aq)$	+2.01					
$H_2O_2(aq) + 2 H^+(aq) + 2 e^- \longrightarrow 2 H_2O(l)$	+1.763					
$MnO_4^{-}(aq) + 8 H^{+}(aq) + 5 e^{-} \longrightarrow Mn^{2+}(aq) + 4 H_2O(l)$	+1.51					
$PbO_2(s) + 4 H^+(aq) + 2 e^- \longrightarrow Pb^{2+}(aq) + 2 H_2O(l)$	+1.455					
$\operatorname{Cl}_2(g) + 2 e^- \longrightarrow 2 \operatorname{Cl}^-(aq)$	+1.358					
$Cr_2O_7^{2-}(aq) + 14 H^+(aq) + 6 e^- \longrightarrow 2 Cr^{3+}(aq) + 7 H_2O(l)$	+1.33					
$MnO_2(s) + 4 H^+(aq) + 2 e^- \longrightarrow Mn^{2+}(aq) + 2 H_2O(l)$	+1.23					
$O_2(g) + 4 H^+(aq) + 4 e^- \longrightarrow 2 H_2O(l)$	+1.229					
$2 \operatorname{IO}_3^{-}(\operatorname{aq}) + 12 \operatorname{H}^+(\operatorname{aq}) + 10 \operatorname{e}^- \longrightarrow \operatorname{I}_2(\operatorname{s}) + 6 \operatorname{H}_2O(\operatorname{l})$	+1.20					
$Br_2(l) + 2 e^- \longrightarrow 2 Br^-(aq)$	+1.065					
$NO_3^{-}(aq) + 4 H^{+}(aq) + 3 e^{-} \longrightarrow NO(g) + 2 H_2O(l)$	+0.956					
$Ag^{+}(aq) + e^{-} \longrightarrow Ag(s)$	+0.800					
$Fe^{3+}(aq) + e^{-} \longrightarrow Fe^{2+}(aq)$	+0.771					
$O_2(g) + 2 H^+(aq) + 2 e^- \longrightarrow H_2O_2(aq)$	+0.695					
$I_2(s) + 2 e^- \longrightarrow 2 I^-(aq)$	+0.535					
$Cu^{2+}(aq) + 2e^{-} \longrightarrow Cu(s)$	+0.340					
$SO_4^{2-}(aq) + 4 H^+(aq) + 2 e^- \longrightarrow 2 H_2O(l) + SO_2(g)$	+0.17					
$\operatorname{Sn}^{4+}(\operatorname{aq}) + 2 \operatorname{e}^{-} \longrightarrow \operatorname{Sn}^{2+}(\operatorname{aq})$	+0.154					
$S(s) + 2 H^+(aq) + 2 e^- \longrightarrow H_2S(g)$	+0.14					
$2 H^{+}(aq) + 2 e^{-} \longrightarrow H_{2}(g)$	0					
$Pb^{2+}(aq) + 2e^{-} \longrightarrow Pb(s)$	-0.125					
$\operatorname{Sn}^{2+}(\operatorname{aq}) + 2 \operatorname{e}^{-} \longrightarrow \operatorname{Sn}(\operatorname{s})$	-0.137					
$Fe^{2+}(aq) + 2e^{-} \longrightarrow Fe(s)$	-0.440					
$Zn^{2+}(aq) + 2e^{-} \longrightarrow Zn(s)$	-0.763					
$Al^{3+}(aq) + 3 e^{-} \longrightarrow Al(s)$	-1.676					
$Mg^{2+}(aq) + 2e^{-} \longrightarrow Mg(s)$	-2.356					
$Na^+(aq) + e^- \longrightarrow Na(s)$	-2.713					
$Ca^{2+}(aq) + 2e^{-} \longrightarrow Ca(s)$	-2.84					
$K^+(aq) + e^- \longrightarrow K(s)$	-2.924					
$Li^+(aq) + e^- \longrightarrow Li(s)$	-3.040					

- 1. You have the following materials from which to construct a galvanic cell: silver metal, magnesium metal, 1.0 M silver (I) nitrate and 1.0 M magnesium nitrate.
 - a. Write the balanced overall reaction that will occur in the galvanic cell. (2 points)
 - b. What electrode will serve as the anode in your galvanic cell? (2 points)
 - c. What is the standard cell potential for your galvanic cell? (4 points)

2. Calculate the time required to produce 2.79 grams if I_2 at an electrode if a current of 1.75 A is passed through a concentrated solution of KI(aq) (8 points)

- 3. Consider an electrochemical cell where the cathode compartment contains Sn(s) and 0.0155 M Sn²⁺(aq) and the anode compartment contains Al(s) and 0.0234 M Al³⁺(aq).
 - a. What is the E° for this electrochemical cell? (3 points)

b. What is the potential for the cell under the conditions given? You may assume a temperature of 298K (6 points)