## JBA 2023 – Chemistry Exam 2

Name:	Coores	/100 =	/00
Name:	Score:	_/100 =	_/00

## Multiple choice questions are worth two points each.

- 1. Constitutional isomers are compounds that have
  - a. the same chemical formulas and molecular structures but different physical properties.
  - b. the same chemical formulas but different molecular structures and physical properties.
  - c. different chemical formulas and molecular structures but the same physical properties.
  - d. the same chemical formulas, molecular structures and physical properties.
- 2. The segment of a polymer shown below represents a

- a. polyester
- b. polyamide.

- c. polyethylene.
- d. polystyrene.

Answer \_\_\_b\_\_\_

Answer b

3. Lidocaine, shown below, is both a local anesthetic and an antiarrhythmic drug. In emergency medical situations, patients with irregular heartbeats frequently receive lidocaine injections or drips. What is the molecular formula of lidocaine?

- a.  $C_{14}H_{20}N_2O$
- b.  $C_{14}H_{17}N_2O$
- c. C<sub>14</sub>H<sub>13</sub>N<sub>2</sub>O
- d.  $C_{14}H_{22}N_2O$

- Answer \_\_d\_\_\_
- 4. Which of the following contains the greatest number of moles of O?
  - a. 2.3 mol H<sub>2</sub>O
- c. 0.9 mol NaNO<sub>3</sub>
- b. 1.2 mol H<sub>2</sub>O<sub>2</sub>
- d. 0.5 mol Ca(NO<sub>3</sub>)<sub>2</sub>
- Answer d

5.	Consider the reaction below.	If 2.00 mol of A reacts with 3.00 mol B, what is the theoretical
	yield of C?	

 $5A + 4B \rightarrow 3C$ 

a. 1.20 mol

c. 3.00 mol

b. 2.25 mol

d. 3.45 mol

Answer a

- 6. One can learn about the environments of hydrogens or carbons present in an organic compound by using the following technique.
  - a. infrared spectroscopy
  - b. mass spectrometry
  - c. NMR spectroscopy
  - d. a really big magnifying glass.

Answer \_\_c\_\_\_

- d. a really big magnifying glass.
- 7. Amino acids are compounds that contain **both** amine and carboxylic acid groups. Which compound is an amino acid? is:

a.

Answer \_\_d\_\_\_

- 8. Which of these classes of compounds form most of the membranes for your cells?
  - a. carbohydrates
  - b. lipids (or fats)
  - c. amino acids

Answer b

- d. nucleic acids.
- 9. What key role do carbohydrates serve in the body?
  - a. They form cell walls
  - b. They encode genetic information
  - c. They are sources of energy

Answer \_\_\_c\_\_

d. They serve no role in the body.

10. Match the term with its definition. (10 points)

10. Water the term with its definition. (10 points)			
Galkene	A. a compound with substituents on the same side of a double bond		
Istereoisomers	B. a collection of atoms in a molecule that has predictable properties and reactivity		
Fchiral carbon	C. a bond that ties two polymer chains together		
_H_mole	D. species that determines the theoretical yield in a reaction		
Ksaturated fat	E. a reactive species with one unpaired electron		
Eradical	F. an atom bonded to 4 different groups		
L_amphiphillic	G. a compound of carbon and hydrogen containing at least one carbon-carbon double bond		
Ccrosslink	H. one Avogadro's number of anything		
Dlimiting reactant	I. compounds that have the same chemical formula and bonding but different arrangement in space		
Bfunctional group	J. a compound of carbon and hydrogen containing at least one carbon-carbon triple bond		
	K. a water insoluble compound with no carbon-carbon multiple bonds		
	L. a containing both hydrophobic and hydrophyllic poritons		

## 11. Complete the following table. (6 points)

12147_ g H <sub>2</sub> O	=	8.14 mol H <sub>2</sub> O	=	4.90 x 10 <sup>24</sup> molecules H <sub>2</sub> O
---------------------------	---	---------------------------	---	--

In order to relate moles and mass, we need to use the molar mass, so we must calculate the molar mass of water:

$$\frac{2 \text{ mol H}}{1 \text{ mol H}_2O}$$
 x  $\frac{1.01 \text{ g}}{1 \text{ mol H}}$  +  $\frac{1 \text{ mol O}}{1 \text{ mol H}_2O}$  x  $\frac{16.00 \text{ g}}{1 \text{ mol O}}$  =  $\frac{18.02 \text{ g}}{1 \text{ mol H}_2O}$ 

Therefore, the molar mass of H<sub>2</sub>O is 18.02 g/mol

Now the conversion between moles and grams:

$$8.14 \frac{\text{mol H}_2 \Theta}{\text{mol H}_2 \Theta} \times \underbrace{18.02 \text{ g}}_{1 \frac{\text{mol H}_2 \Theta}{\text{mol H}_2 \Theta}} = 146.68 \text{ g} = 147 \text{ g H}_2 \Theta$$

To convert between moles and molecules, we use Avogadro's number that tells us that  $1 \text{ mol} = 6.022 \times 10^{23} \text{ molecules}$ .

10. Identify the functional group shown in each structure: (4 points)

- 11. Kevlar, a poly**amide** used to make bulletproof vests, is made from terephthalic acid and paraphenylenediamine.
- a. Write the polymerization reaction for the formation of Kevlar, indicating the repeating structure for the polymer. (4 points)

b. What type of polymerization reaction is this? Circle the correct answer. (2 points)

addition condensation

12. Complete the table for the alkanes below. (6 points)

Structure (Line angle or Lewis structures are acceptable)	Name
	4-ethyl-6-methyl nonane
H <sub>3</sub> C <sub>CH</sub> CH <sub>3</sub>     CH <sub>2</sub>   H <sub>3</sub> C-CH <sub>2</sub> -C-CH <sub>3</sub>   CH <sub>3</sub>	2,4,4-trimethyhexane

- 13. Draw organic compounds that fit the following criteria (there may be more than one structure that fits the criteria, you only need to draw one example for each):
  - a. A compound that contains an ester and has the formula  $C_5H_{10}O_2$ . (4 points)

Here are examples of possible structures:

b. A compound that contains and amine and an ether and has the formula  $C_4H_{11}NO$ . (4 points)

Here's an example of one possible structure:

14. Compounds that can serve as monomers for polymerization reactions must have one key property. What property is this? Show how this property appears in both addition and condensation polymerization. (6 points)

Monomers must be able to react in two locations in order for the polymer to continue to grow. For addition polymerization, the alkene produces a di-radical that can react in two locations and continue to grow. For condensation polymerization, the monomers must have two functional groups that can react independently. (Example structures would be useful here.)

15. Complete the following table (12 points)

Line angle	Lewis Structure	Molecular Formula
НО	H C H H H H	C <sub>5</sub> H <sub>10</sub> O <sub>2</sub>
N	HC H H H H H H H H H H H H H H H H H H	C <sub>9</sub> H <sub>13</sub> N
НО	H — C — H	C5H8O

16. For the reaction below, what is the limiting reactant when 0.253 g aluminum reacts with 0.482 g Cl<sub>2</sub>? You must justify your answer with a calculation. (8 pts)  $2Al(s) + 3Cl_2(g) \rightarrow 2AlCl_3(s)$ 

If aluminum is the L.R. how much AlCl<sub>3</sub> could be made?

$$0.253 \ \frac{\text{g Al}}{\text{g Al}} \ \ \text{x} \ \ \frac{1 \ \text{mol Al}}{26.98 \ \frac{\text{g Al}}{\text{g Al}}} \ \ \text{x} \ \ \frac{2 \ \text{mol AlCl}_3}{2 \ \text{mol Al}} = \ \ \textbf{0.00938 \ mol AlCl}_3 \ \ \text{x} \ \ \frac{133.34 \ \text{g AlCl}_3}{1 \ \text{mol AlCl}_3} = \ \ \textbf{1.25g AlCl}_3$$

If chlorine is the L.R. how much AlCl<sub>3</sub> could be made?

$$0.482 \ \frac{\text{g-Cl}_2}{70.90} \ x \ \underline{\frac{1 \ \text{mol Cl}_2}{70.90}} \ x \ \underline{\frac{2 \ \text{mol AlCl}_3}{3 \ \text{mol Cl}_2}} = \ \textbf{0.00453 \ mol AlCl}_3 \ x \ \underline{\frac{133.34 \ \text{g AlCl}_3}{1 \ \text{mol AlCl}_3}} = \ \textbf{0.604g \ AlCl}_3$$

Since less AlCl<sub>3</sub> is produced when all of the Cl<sub>2</sub> is consumed, Cl<sub>2</sub> must be the limiting reagent.

17. Alcohols and carboxylic acids react to form esters by the process shown below.

a. Complete the reaction of salicylic acid with methanol to form methyl salicylate (oil of wintergreen). (4 points)

b. Complete the reaction of salicylic acid with acetic acid to form acetylsalicylic acid (aspirin). (4 points)

- 13. Nucleic acids are the building blocks of DNA.
  - a. Every nucleic acid in DNA has three components. What are they? (3 points)

A sugar, a phosphate group and a nitrogen containing base

b. Why types of intermolecular forces hold DNA strands together to form a double helix? (3 points)

## Hydrogen bonding interactions

c. In DNA, adenine and thymine are complementary bases, as are cytosine and guanine. If a segment of a single strand of DNA has the sequence, -ATTCGTAA-, what would the complementary sequence be? (4 points)

Since A is complementary to T and C is complementary to G, the sequence would be:
-TAAGCATT-